

All Databases PubMed Nucleotide Protein Genome Structure OMIM PACC
Journals BooksSearch PubMed for [Advanced Search \(beta\)](#)[Limits](#) [Preview/Index](#) [History](#) [Clipboard](#) [Details](#)

Display AbstractPlus Show 20 Sort By Send to

All: 1 Review: 0

☐ 1: [Anticancer Res. 1992 Jan-Feb;12\(1\):21-5.](#)[Links](#)**Modulation of multidrug resistance by immunosuppressive agents: cyclosporin analogues, FK506 and mizoribine.**[Mizuno K](#), [Furuhashi Y](#), [Misawa T](#), [Iwata M](#), [Kawai M](#), [Kikkawa F](#), [Kano T](#), [Tomoda Y](#).

Department of Obstetrics and Gynecology, Nagoya University School of Medicine, Japan.

It has recently been reported that an immunosuppressive agent, cyclosporin A, shows a potent overcoming effect on multidrug resistance (MDR). We studied the presence of such a modulating effect of cyclosporin analogues and other immunosuppressive agents, FK506 and mizoribine, in human multidrug-resistant ovarian cancer cells (TAOV/A0.2). The intensity of the overcoming effect of cyclosporin analogues against adriamycin resistance was found to be in the order of cyclosporin D greater than A greater than C greater than H. It was found that cyclosporin D, which has relatively weak immunosuppressive activity, overcame adriamycin resistance in the multidrug-resistant ovarian cancer cells to a remarkable degree. On the other hand, it was found that FK506, a new potent immunosuppressant, could also distinctly modulate adriamycin-resistance. It was found that FK506 conferred chemosensitization upon adriamycin with reincreasing intracellular adriamycin accumulation in MDR cells which was far less than the parent strain. However, in the case of mizoribine, no modulation of drug resistance existed. Such modulation was not necessarily accompanied by immunosuppressive activity and the two functions were thought to be based on different mechanisms.

PMID: 1373592 [PubMed - indexed for MEDLINE]

Related Articles

Secondary combined resistance to the multidrug-resistance-reversing activity of cyclosporin A in the cell line F4-6RADR-CsA. [*J Cancer Res Clin Oncol.* 1994]

Modulation of adriamycin cytotoxicity and transport in drug-sensitive and multidrug-resistant Chinese hamster ovary cells by hypochlorous acid and sodium iodide. [*Chem Biol Interact.* 2000]

Reversal of multidrug resistance in Chinese hamster ovary cells by the immunosuppressant cyclosporin A. [*Int J Cancer.* 1993]

Intracellular levels of two cyclosporin derivatives valspodar (PSC 833) and cyclosporin A closely associated with multidrug resistance-modulating activity in sublines of human colorectal adenocarcinoma. [*J Cancer Res.* 2001]

FK506 reverses adriamycin resistance in a multidrug-resistant human leukemia cell line. [*Kobe J Med Sci.* 1992]

[» See all Related Articles...](#)

Patient Drug Information

Daunorubicin (Cerubidine®, DaunoXome®) Your doctor has ordered the drug daunorubicin to help treat your illness. The drug is given by injection into a vein.

[» read more ...](#)

Doxorubicin (Adriamycin PFS®, Adriamycin RDP®, Adriamycin®, ...) Your doctor has ordered the drug doxorubicin to help treat your illness. The drug is given by injection into a vein.

[» read more ...](#)

Cyclosporine (Neoral[®], Sandimmune[®], Gengraf[®]) Cyclosporine and cyclosporine (modified) are used with other medications to prevent transplant rejection (attack of the transplanted organ by the immune system of the person who received the organ) in people who have rec...

» read more ...

» See all 4 Drug Reports ...

Display Show Sort By Send to

[Write to the Help Desk](#)

[NCBI](#) | [NLM](#) | [NIH](#)

[Department of Health & Human Services](#)

[Privacy Statement](#) | [Freedom of Information Act](#) | [Disclaimer](#)